

# On Testing Convexity and Submodularity

Michal Parnas  
The Academic College  
of Tel-Aviv-Yaffo  
Tel-Aviv, ISRAEL  
michalp@mta.ac.il

Dana Ron  
Department of EE – Systems  
Tel-Aviv University  
Ramat Aviv, ISRAEL  
danar@eng.tau.ac.il

Ronitt Rubinfeld  
NEC Research Institute  
Princeton, NJ  
ronitt@research.nj.nec.com

September 4, 2002

## Abstract

Convex and Submodular functions play an important role in many applications, and in particular in combinatorial optimization. Here we study two special cases: convexity in one dimension and submodularity in two dimensions. The latter type of functions are equivalent to the well known *Monge matrices*. A matrix  $V = \{v_{i,j}\}_{i,j=0}^{i=n_1, j=n_2}$  is called a Monge matrix if for every  $0 \leq i < i' \leq n_1$  and  $0 \leq j < j' \leq n_2$ , we have  $v_{i,j} + v_{i',j'} \leq v_{i,j'} + v_{i',j}$ . If inequality holds in the opposite direction then  $V$  is an *inverse Monge* matrix (supermodular function). Many problems, such as the traveling salesperson problem and various transportation problems, can be solved more efficiently if the input is a Monge matrix.

In this work we present testing algorithms for the above properties. A Testing algorithm for a predetermined property  $\mathcal{P}$  is given query access to an unknown function  $f$ , and a distance parameter  $\epsilon$ . The algorithm should accept  $f$  with high probability if it has the property  $\mathcal{P}$ , and reject it with high probability if more than an  $\epsilon$ -fraction of the function values should be modified so that  $f$  obtains the property. Our algorithm for testing whether a one-dimensional function  $f : [n] \rightarrow \mathbb{R}$  is convex (concave), has query complexity and running time of  $O((\log n)/\epsilon)$ . Our algorithm for testing whether an  $n_1 \times n_2$  matrix  $V$  is a Monge (inverse Monge) matrix has query complexity and running time of  $O((\log n_1 \cdot \log n_2)/\epsilon)$ .

# 1 Introduction

Convex functions and their combinatorial analogs, submodular functions, play an important role in many disciplines and applications, including combinatorial optimization, game theory, probability theory, and electronic trade. Such functions exhibit a rich mathematical structure (see Lovász [Lov83]), which often makes it possible to efficiently find their minimum [GLS81, IFF00, Sch00], and thus leads to efficient algorithms for many important optimization problems.

Convex functions over discrete domains are defined as follows.

**Definition 1 (Convex and Concave)** *Let  $f$  be a function defined over a discrete domain  $X$ . The function  $f$  is convex if for all  $x, y \in X$  and for all  $0 \leq \alpha \leq 1$  such that  $\alpha x + (1 - \alpha)y \in X$ , it holds that  $f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y)$ . The function  $f$  is concave if for all  $x, y \in X$  and for all  $0 \leq \alpha \leq 1$  such that  $\alpha x + (1 - \alpha)y \in X$ , it holds that  $f(\alpha x + (1 - \alpha)y) \geq \alpha f(x) + (1 - \alpha)f(y)$ .*

Submodular functions are defined as follows: Let  $\mathcal{I} = I_1 \times I_2 \times \dots \times I_d$ ,  $d \geq 2$ , be a product space where  $I_q \subseteq \mathbb{R}$ . In particular, we are interested in discrete domains  $I_q = \{0, \dots, n_q\}$ . The *join* and *meet* operations are defined for every  $x, y \in \mathcal{I}$ :

$$(x_1, \dots, x_d) \vee (y_1, \dots, y_d) \stackrel{\text{def}}{=} (\max\{x_1, y_1\}, \dots, \max\{x_d, y_d\})$$

and

$$(x_1, \dots, x_d) \wedge (y_1, \dots, y_d) \stackrel{\text{def}}{=} (\min\{x_1, y_1\}, \dots, \min\{x_d, y_d\}),$$

respectively.

**Definition 2 (Submodularity and Supermodularity)** *A function  $f : \mathcal{I} \rightarrow \mathbb{R}$  is submodular if for every  $x, y \in \mathcal{I}$ ,  $f(x \vee y) + f(x \wedge y) \leq f(x) + f(y)$ . The function  $f$  is supermodular if for every  $x, y \in \mathcal{I}$ ,  $f(x \vee y) + f(x \wedge y) \geq f(x) + f(y)$ .*

Certain subclasses of submodular functions are of particular interest. One such subclass is that of *submodular set* functions, which are defined over binary domains. That is,  $I_q = \{0, 1\}$  for every  $1 \leq q \leq d$ , and so each  $x \in \mathcal{I}$  corresponds to a subset of  $\{1, \dots, d\}$ . Such functions are used for example in the scenario of combinatorial auctions on the internet (e.g. [dVV00],[LLN01]).

Another important subclass is the class of *Monge* functions, which are obtained when the domain is large but the dimension is  $d = 2$ . Since such functions are 2-dimensional, it is convenient to represent them as 2-dimensional matrices, which are referred to as *Monge matrices*. When the function is a 2-dimensional supermodular function the corresponding matrix is called an *inverse Monge matrix*.

The first problem that was shown to be solvable more efficiently if the underlying cost matrix is a Monge matrix is the classical Hitchcock transportation problem (see Hoffman [Hof63]). Since then it has been shown that many other combinatorial optimization problems can be solved more efficiently in this case (e.g. weighted bipartite matching, and NP-hard problems such as the traveling salesperson problem). See [BKR96] for a comprehensive survey on Monge matrices and their applications.

## 1.1 Testing Convexity and Submodularity

In this paper we approach the questions of convexity and submodularity from within the framework of property testing [RS96, GGR98]. (For surveys on property testing see [Ron01, Fis01].) Let  $f$  be a fixed but unknown function, and let  $\mathcal{P}$  be a fixed property of functions (such as the convexity or submodularity of a function). A testing algorithm for the property  $\mathcal{P}$  should determine, by querying  $f$ , whether  $f$  has the property  $\mathcal{P}$ , or whether it is  $\epsilon$ -far from having the property for a given distance parameter  $\epsilon$ . By  $\epsilon$ -far we mean that more than an  $\epsilon$ -fraction of the values of  $f$  should be modified so that  $f$  obtains the desired property  $\mathcal{P}$ .

**Our Results.** We present efficient testing algorithms for discrete convexity in one dimension and for Monge matrices. Specifically:

- We describe and analyze an algorithm that tests whether a function  $f : [n] \rightarrow \mathbb{R}$  is convex (concave). The running time of this algorithm is  $O(\log n/\epsilon)$ .
- We describe and analyze a testing algorithm for Monge and inverse Monge matrices whose running time is  $O((\log n_1 \cdot \log n_2)/\epsilon)$ , when given an  $n_1 \times n_2$  matrix.

Furthermore, the testing algorithm for inverse Monge matrices can be used to derive a testing algorithm, with the same complexity, for an important sub-family of Monge matrices, named *distribution matrices*. A matrix  $V = \{v_{i,j}\}$  is said to be a distribution matrix, if there exists a non-negative *density matrix*  $D = \{d_{k,\ell}\}$ , such that every entry  $v_{i,j}$  in  $V$  is of the form  $v_{i,j} = \sum_{k \leq i} \sum_{\ell \leq j} d_{k,\ell}$ . In other words, the entry  $v_{i,j}$  corresponds to the cumulative density of all entries  $d_{k,\ell}$  such that  $k \leq i$  and  $\ell \leq j$ .

Thus in both cases the complexity of the algorithms is linear in  $1/\epsilon$  and polylogarithmic in the size of the domain.

## 1.2 Techniques.

**Convexity in One Dimension.** We start with the following basic observation: A function  $f : [n] \rightarrow \mathbb{R}$  is convex if and only if for every  $1 \leq i \leq n-1$ ,  $(f(i+1) - f(i)) - (f(i) - f(i-1)) \geq 0$ . Given this characterization, consider the *difference* function  $f'$  which is defined as  $f'(i) = f(i) - f(i-1)$ . That is,  $f'$  is a discrete analog of the first derivative of  $f$ . By the above observation we have that  $f$  is convex if and only if  $f'$  is monotone non-decreasing. Hence, a tempting approach for testing whether  $f$  is convex would be to test whether  $f'$  is monotone non-decreasing, where this can be done in time  $O(\log^2 n/\epsilon)$  [EKK<sup>+</sup>00, BRW99, DGL<sup>+</sup>99].

Unfortunately this approach does not work. There are functions  $f$  that are very far from convex but their difference function  $f'$  is very close to monotone.<sup>1</sup> Therefore, instead of considering only consecutive points  $i, i+1$ , we consider pairs of points  $i, j \in [n]$  that are not necessarily consecutive. More precisely, we select intervals  $\{i, \dots, j\}$  of varying lengths and check that for each interval selected, certain constraints are satisfied. If  $f$  is convex then these constraints are satisfied for every interval. On the other hand, we show that if  $f$  is  $\epsilon$ -far from convex then the probability that we observe a violation of some constraint is sufficiently large.

**Monge Matrices.** As stated above, it is convenient to represent 2-dimensional submodular functions as 2-dimensional Monge matrices. Thus a function  $f : \{0, \dots, n_1\} \times \{0, \dots, n_2\} \rightarrow \mathbb{R}$  can be

---

<sup>1</sup>In particular consider the function  $f$  such that for every  $i \leq n/2$ ,  $f(i) = i$ , and for  $i > n/2$ ,  $f(i) = i - 1$ . In other words,  $f'(i) = 1$  for every  $i$  except  $i = n/2$  where  $f'(i) = 0$ . Then  $f'$  is very close to monotone, but it is not hard to verify that  $f$  is far from convex.

represented as the matrix  $V = \{v_{i,j}\}_{i,j=0}^{i=n_1, j=n_2}$  where  $v_{i,j} = f(i, j)$ . Observe that for every pair of indices  $(i, j'), (i', j)$  such that  $i < i'$  and  $j < j'$  we have that  $(i, j') \vee (i', j) = (i', j')$  and  $(i, j') \wedge (i', j) = (i, j)$ . It follows from Definition 2 that  $V$  is a Monge matrix ( $f$  is a 2-dimensional submodular function) if and only if:

$$\forall i, j, i', j' \text{ s.t. } i < i', j < j' : \quad v_{i,j} + v_{i',j'} \leq v_{i,j'} + v_{i',j}$$

and  $V$  is an inverse Monge matrix ( $f$  is a 2-dimensional supermodular function) if and only if:

$$\forall i, j, i', j' \text{ s.t. } i < i', j < j' : \quad v_{i,j} + v_{i',j'} \geq v_{i,j'} + v_{i',j} .$$

That is, in both cases we have a constraint for every quadruple  $v_{i,j}, v_{i',j'}, v_{i,j'}, v_{i',j}$  such that  $i < i'$  and  $j < j'$ .<sup>2</sup> Our algorithm selects such quadruples according to a particular (non-uniform) distribution and verifies that the constraint is satisfied for every quadruple selected. Clearly the algorithm always accepts Monge matrices. The main thrust of the analysis is in showing that if the matrix  $V$  is far from being Monge then the probability of obtaining a “bad” quadruple is sufficiently large.

A central building block in proving the above, is the following combinatorial problem, which may be of independent interest. Let  $C$  be a given matrix, possibly containing negative values, and let  $R$  be a subset of positions in  $C$ . We are interested in refilling the entries of  $C$  that reside in  $R$  with *non-negative* values, such that the following constraint is satisfied: for every position  $(i, j)$  that does not belong to  $R$ , the sum of the modified values in  $C$  that are below<sup>3</sup>  $(i, j)$ , is the same as in the original matrix  $C$ . That is, the sum of the modified values in entries  $(k, \ell)$ , such that  $k \leq i$  and  $j \leq \ell$ , remains as it was.

We provide sufficient conditions on  $C$  and  $R$  under which the above is possible, and describe the corresponding procedure that refills the entries of  $C$  that reside in  $R$ . Our starting point is a simple special case in which  $R$  corresponds to a sub-matrix of  $C$ . In such a case it suffices that for each row and each column in  $R$ , the sum of the corresponding entries in the original matrix  $C$  is non-negative. Under these conditions a simple greedy algorithm can modify  $C$  as required. Our procedure for general subsets  $R$  is more involved but uses the sub-matrix case as a subroutine.

### 1.3 Further Research

We suggest the following open problems. First it remains open to determine the complexity of testing discrete convexity (concavity) when the dimension  $d$  of the input domain is greater than 1, and for testing submodular (supermodular) functions when the dimension  $d$  is greater than 2. Note that though submodular functions can be viewed as a certain interpretation of convexity in dimensions  $d \geq 2$ , they do not necessarily satisfy Definition 1.

It seems that our algorithm for testing Monge matrices and its analysis can be extended to work for testing the special case of distribution matrices of dimension  $d > 2$ , where the complexity of the resulting algorithm is  $O\left(\left(\prod_{q=1}^d \log n_q\right)/\epsilon\right)$ . However, as opposed to the  $d = 2$  case, where Monge matrices are only slightly more general than distribution matrices, for  $d > 2$  Monge matrices are more expressive. Hence it is not immediately clear how to adapt our algorithm to testing Monge matrices in higher dimensions.

<sup>2</sup>It is easy to verify that for all other  $i, j, i', j'$  (with the exception of the symmetric case where  $i' < i$  and  $j' < j$ ), the constraint holds trivially (with equality).

<sup>3</sup>We denote the lower left position of the matrix  $C$  by  $(0, 0)$ .

It would also be interesting to find an efficient testing algorithm for the subclass of submodular set functions, which are defined over binary domains.

Finally, in many optimization problems it is enough that the underlying cost matrix is a permutation of a Monge matrix. In such cases it may be useful to test whether a given matrix is a permutation of some Monge matrix or far from any permuted Monge matrix.

**Organization.** The testing algorithm for convexity is described in Section 2. The remainder of the paper is dedicated to testing Monge matrices. In Section 3 we describe several building blocks that will be used by our testing algorithm for Monge matrices. In Section 4 we describe a testing algorithm for Monge matrices whose complexity is  $O(n/\epsilon)$ , where we assume for simplicity that the matrix is  $n \times n$ . Building on this algorithm and its analysis, in Section 5 we present a significantly faster algorithm whose complexity is  $O((\log^2 n)/\epsilon)$ . We conclude this section with a short discussion concerning distribution matrices.

## 2 Testing Convexity in 1-Dimension

As noted in the introduction, in the case that the domain is  $X = [n] = \{0, \dots, n\}$ , we get the following characterization for convexity, whose proof is included for completeness.

**Claim 1 (1-D Convex)** *A function  $f : [n] \rightarrow \mathbb{R}$  is convex if and only if for all  $1 \leq i \leq n - 1$ ,  $f(i) - f(i - 1) \leq f(i + 1) - f(i)$ .*

**Proof:** If  $f$  is convex then in particular for  $x = i - 1$ ,  $y = i + 1$  and  $\alpha = 1/2$  we have  $\alpha x + (1 - \alpha)y = \frac{i-1}{2} + \frac{i+1}{2} = i$ . By Definition 1,  $f(i) \leq \frac{1}{2}f(i - 1) + \frac{1}{2}f(i + 1)$ , or equivalently,  $f(i) - f(i - 1) \leq f(i + 1) - f(i)$ .

In the other direction, suppose that  $f(i) - f(i - 1) \leq f(i + 1) - f(i)$  for every  $1 \leq i \leq n - 1$ . Consider any  $x, y \in [n]$  and  $0 < \alpha < 1$  such that  $z = \alpha \cdot x + (1 - \alpha) \cdot y$  is an integer. Assume without loss of generality that  $x < y$ . Now we have that

$$f(y) - f(y - 1) \geq f(y - 1) - f(y - 2) \geq \dots \geq f(z + 1) - f(z) \geq f(z) - f(z - 1) \geq \dots \geq f(x + 1) - f(x).$$

Then, since the differences are monotone non-increasing, the average of the first  $\alpha(y - x)$  differences is greater or equal to the average of the next  $(1 - \alpha)(y - x)$  differences. Since  $z = y - \alpha(y - x) = x + (1 - \alpha)(y - x)$ , we have that

$$\frac{(f(y) - f(y - 1)) + (f(y - 1) - f(y - 2)) + \dots + (f(z + 1) - f(z))}{\alpha(y - x)} \tag{1}$$

$$\geq \frac{(f(z) - f(z - 1)) + (f(z - 1) - f(z - 2)) + \dots + (f(x + 1) - f(x))}{(1 - \alpha)(y - x)}. \tag{2}$$

This is equivalent to  $(1 - \alpha)(f(y) - f(z)) \geq \alpha(f(z) - f(x))$ , that is  $f(z) \leq \alpha f(x) + (1 - \alpha)f(y)$ , as required. ■

Denote by  $I_{i,j}$  the interval  $\{i, i + 1, \dots, j\}$  of points. Let  $mid = \lfloor (i + j)/2 \rfloor$  be the mid point of  $I_{i,j}$ .

**Definition 3** *For every  $0 \leq i < j \leq n$  such that  $j - i > 7$ , we say that the interval  $I_{i,j}$  is good with respect to  $f$  if the following holds:*

$$\begin{aligned} f(i + 1) - f(i) &\leq \frac{f(mid - 1) - f(i + 1)}{(mid - 1) - (i + 1)} \leq f(mid) - f(mid - 1) \leq f(mid + 1) - f(mid) \\ &\leq f(mid + 2) - f(mid + 1) \leq \frac{f(j - 1) - f(mid + 2)}{(j - 1) - (mid + 2)} \leq f(j) - f(j - 1) \end{aligned}$$

Otherwise we say that the interval is bad with respect to  $f$ . If  $j - i \leq 7$ , then  $I_{i,j}$  is good with respect to  $f$  if and only if the function  $f$  is convex over  $I_{i,j}$ .

In order to test if  $f$  is convex we test recursively if sub-intervals of  $I_{0,n}$  are good.

**Algorithm 1 Test-Convex**

1. Repeat  $2/\epsilon$  times: Test-Interval( $I_{0,n}$ ).
2. If all of the tests in Step 1 accepted then accept, otherwise reject.

**Procedure Test-Interval( $I_{i,j}$ )**

1. Check that  $I_{i,j}$  is good with respect to  $f$ . In not, reject.
2. If  $j - i > 7$  then: Uniformly at random call either Test-Interval( $I_{i,mid}$ ) or Test-Interval( $I_{mid+1,j}$ ), where  $mid = \lfloor (i + j)/2 \rfloor$ .
3. If the test in Step 2 accepted then accept, otherwise reject.

**Theorem 1** *If  $f$  is convex then Algorithm 1 always accepts, and if  $f$  is  $\epsilon$ -far from convex then the algorithm rejects with a probability of at least  $2/3$ .*

**Proof:** For the sake of brevity, unless stated otherwise, when we say that an interval is good, then we mean with respect to  $f$ . If  $f$  is convex then all intervals  $I_{i,j}$  are good, and hence Algorithm 1 accepts with probability 1. In order to prove that if  $f$  is  $\epsilon$ -far from convex then the algorithm rejects with probability of at least  $2/3$ , we prove the contrapositive statement. Assume that the algorithm accepts with a probability greater than a  $1/3$ . We will show that  $f$  is  $\epsilon$ -close to a convex function.

To this end we define a tree, whose vertices correspond to all possible intervals  $I_{i,j}$  that may be tested recursively in calls to Test-Interval( $I_{i,j}$ ). Specifically, the root of the tree corresponds to  $I_{0,n}$ . The children of the internal vertex corresponding to  $I_{i,j}$  are the vertices corresponding to  $I_{i,mid}$  and  $I_{mid+1,j}$ , where  $mid = \lfloor (i + j)/2 \rfloor$ . The leaves of the tree correspond to the smallest intervals tested, that is, intervals  $I_{i,j}$  for which  $j - i \leq 7$ .

We say that an internal vertex in the tree is good if the corresponding interval is good. We say that a leaf is good if its interval and all its ancestors are good. Otherwise, the vertex (leaf) is bad. We say that a path from the root to a leaf is good if all vertices along it are good. Otherwise the path is bad. For each level  $\ell$  in the tree,  $\ell = 0, \dots, \log n$ , let  $\mathcal{B}_\ell$  be the subset of vertices in the the  $\ell$ 'th level of the tree that are bad but whose ancestors are all good. Let  $\mathcal{B} = \bigcup_\ell \mathcal{B}_\ell$ , and let  $\epsilon_\ell$  be the fraction of vertices in level  $\ell$  of the tree that belong to  $\mathcal{B}_\ell$ .

**Sub-Claim 1** *If Algorithm 1 accepts  $f$  with a probability greater than a  $1/3$ , then  $\sum_\ell \epsilon_\ell \leq \epsilon$ .*

**Proof:** Assume by contradiction that  $\sum_\ell \epsilon_\ell > \epsilon$ . Observe that by the definition of  $\mathcal{B}$ , all leaves which are descendents of a vertex in  $\mathcal{B}$  are bad, and every bad leaf either belongs to  $\mathcal{B}$  or has a single ancestor in  $\mathcal{B}$ . Therefore, if  $\sum_\ell \epsilon_\ell > \epsilon$ , then the fraction of bad leaves is greater than  $\epsilon$ . But in such a case, the probability that the algorithm does not follow a bad path to a bad leaf (passing through a vertex in  $\mathcal{B}$ ), in any one of its  $2/\epsilon$  iterations, is at most  $(1 - \epsilon)^{2/\epsilon} < e^{-2} < 1/3$ . This contradicts our assumption that the algorithm accepts with a probability greater than a  $1/3$ .  $\square$

Hence we assume from now on that  $\sum_{\ell} \epsilon_{\ell} \leq \epsilon$ . We show how to modify  $f$  in at most  $\epsilon \cdot n$  places so that the resulting function, denoted  $g$ , is convex. In particular, we shall modify the value of  $f$  on every bad interval  $I_{i,j}$  whose corresponding vertex in the tree belongs to  $\mathcal{B}$ . The value of  $g$  is defined to be the same as the value of  $f$  on all points outside of these intervals. Since  $\sum_{\ell} \epsilon_{\ell} \leq \epsilon$ , the total fraction of points modified is at most  $\epsilon$  as required. Observe that by the definition of the tree and  $\mathcal{B}$ , for every two intervals whose corresponding vertices belong to  $\mathcal{B}$ , the intersection of the intervals is empty. Hence we can modify each one of these intervals independently.

Let  $I_{i,j}$  be a bad interval corresponding to a vertex in  $\mathcal{B}$ . We modify  $I_{i,j}$  as follows:

- $f(i), f(i+1), f(j-1)$  and  $f(j)$  remain unchanged. That is, set  $g(i) = f(i), g(i+1) = f(i+1), g(j-1) = f(j-1)$  and  $g(j) = f(j)$ .
- For every  $t, i+1 < t < j-1$ , set  $g(t) = f(i+1) + \frac{f(j-1)-f(i+1)}{(j-1)-(i+1)} \cdot (t - (i+1))$ .

*Sub-Claim 2 The function  $g$  is convex.*

*Proof:* We shall show first that all intervals  $I_{i,j}$  corresponding to vertices in the tree are good with respect to  $g$ , and from this derive the convexity of  $g$ .

We start with the first part. Consider any such interval  $I_{i,j}$  whose corresponding vertex in the tree is  $v$ . Let  $Anchor = \{i, i+1, mid-1, mid, mid+1, mid+2, j-1, j\}$  be the set of points which participate in the definition of a good interval  $I_{i,j}$ . We will show that the value of  $g$  on points  $p \in Anchor$  is such that the interval  $I_{i,j}$  is good with respect to  $g$ . There are two cases:

1. The interval  $I_{i,j}$  is good with respect to  $f$ , and  $v$  does not have any ancestors in  $\mathcal{B}$ . If  $v$  also has no descendents in  $\mathcal{B}$ , then it clearly remains good with respect to  $g$ , since no modification is performed on any point in the interval, and so  $g(t) = f(t)$  for every  $i \leq t \leq j$ . Otherwise,  $v$  has a descendent in  $\mathcal{B}$ . In this case, let  $p \in Anchor$ , let  $v'$  be a descendent of  $v$ , and let  $I_{i',j'}$  denote the interval corresponding to  $v'$ . If  $i' \leq p \leq j'$ , then by definition of the tree, either  $p = i'$  or  $p = i'+1$  or  $p = j'-1$  or  $p = j'$ . Therefore, even if  $v' \in \mathcal{B}$  and the interval  $I_{i',j'}$  is modified, then by the definition of  $g$  we have that  $g(p) = f(p)$  for every  $p \in Anchor$ . Thus  $I_{i,j}$  remains good with respect to  $g$ .
2. Either  $v \in \mathcal{B}$  or  $v$  has an ancestor  $v' \in \mathcal{B}$ . In this case let  $I_{i',j'}$  be the corresponding interval of  $v'$ . By definition,  $I_{i,j} \subseteq I_{i',j'}$ . Since the parent of  $I_{i',j'}$  is good, we know that

$$f(i'+1) - f(i') \leq \frac{f(j'-1) - f(i'+1)}{(j'-1) - (i'+1)} \leq f(j') - f(j'-1).$$

It is not hard to verify that by the definition of  $g$  we get that  $g(t) - g(t-1) \leq g(t+1) - g(t)$  for every  $i' < t < j'$ , and in particular for every  $i < t < j$ . It follows that  $I_{i,j}$  is good with respect to  $g$ .

Hence all intervals corresponding to vertices in the tree are good with respect to  $g$ . We now prove that for every  $0 < t < n$  it holds that  $g(t) - g(t-1) \leq g(t+1) - g(t)$ , and thus  $g$  is convex. Let  $I_{i,j}$  be the smallest interval in the tree such that  $i < t < j$ . If  $j - i \leq 7$  then we are done, since the goodness of  $I_{i,j}$  in this case means that  $g$  is convex over the whole interval. Otherwise, either  $t = mid$  or  $t = mid + 1$ , where  $mid = \lfloor (i+j)/2 \rfloor$ . To verify this, note that if this were not the case then either  $i < t < mid$  or  $mid + 1 < t < j$ . Hence  $t$  is contained in a smaller interval in the tree, contradicting the minimality of  $I_{i,j}$ . But since  $I_{i,j}$  is good with respect to  $g$ ,  $g(mid) - g(mid-1) \leq g(mid+1) - g(mid)$ , and  $g(mid+1) - g(mid) \leq g(mid+2) - g(mid+1)$ . Thus we are done with the proof of Sub-Claim 2, and Theorem 1 follows. ■

### 3 Building Blocks for Our Algorithms for Testing Inverse Monge

From this point on we focus on inverse Monge matrices. Analogous claims hold for Monge matrices. We also assume for simplicity that the dimensions of the matrices are  $n_1 = n_2 = n$ . In what follows we provide a characterization of inverse Monge matrices that is exploited by our algorithms. Given any real valued matrix  $V = \{v_{i,j}\}_{i,j=0}^{i,j=n}$  we define an  $(n+1) \times (n+1)$  matrix  $C'_V = \{c_{i,j}\}_{i,j=0}^{i,j=n}$  as follows:

- $c_{0,0} = v_{0,0}$ ;
- For  $i > 0$ :  $c_{i,0} = v_{i,0} - v_{i-1,0}$ ;
- For  $j > 0$ :  $c_{0,j} = v_{0,j} - v_{0,j-1}$ ;
- And for every  $i, j > 0$ :

$$\begin{aligned} c_{i,j} &= (v_{i,j} - v_{i-1,j}) - (v_{i,j-1} - v_{i-1,j-1}) \\ &= (v_{i,j} - v_{i,j-1}) - (v_{i-1,j} - v_{i-1,j-1}). \end{aligned} \quad (3)$$

Let  $C_V = \{c_{i,j}\}_{i,j=1}^{i,j=n}$  be the sub-matrix of  $C'_V$  that includes all but the first (0'th) row and column of  $C'_V$ . The following two claims are well known and easy to verify. We include their proofs for completeness.

**Claim 2** For every  $0 \leq i, j \leq n$ ,  $v_{i,j} = \sum_{k=0}^i \sum_{\ell=0}^j c_{k,\ell}$ .

**Proof:** The claim is proved by induction on  $i$  and  $j$ .

The base case,  $i, j = 0$  holds by definition of  $c_{0,0}$ .

Consider any  $i > 0$  and assume that the claim holds for every  $k < i$ ,  $j = 0$ . We prove it for  $i$  and for  $j = 0$ . By definition of  $c_{i,0}$  we have  $v_{i,0} = v_{i-1,0} + c_{i,0}$ . By the induction hypothesis,  $v_{i-1,0} = \sum_{k=0}^{i-1} c_{k,0}$ , and the induction step follows. The claim is similarly proved for every  $j > 0$  and  $i = 0$ .

Finally, consider any  $i, j > 0$  and assume that the claim holds for every  $k < i$  and  $\ell \leq j$ , and for every  $k \leq i$  and  $\ell < j$ . We prove it for  $i, j$ . By definition of  $c_{i,j}$ ,  $v_{i,j} = v_{i-1,j} + (v_{i,j-1} - v_{i-1,j-1}) + c_{i,j}$ . By the induction hypothesis,

$$v_{i-1,j} + (v_{i,j-1} - v_{i-1,j-1}) = \sum_{k=0}^{i-1} \sum_{\ell=0}^j c_{k,\ell} + \sum_{\ell=0}^{j-1} c_{i,\ell}$$

and the induction step follows. ■

**Claim 3** A matrix  $V$  is an inverse Monge matrix if and only if  $C_V$  is a non-negative matrix.

**Proof:** If  $V$  is an inverse Monge matrix, then in particular, for every  $i, j \geq 1$  we have that  $v_{i,j} + v_{i-1,j-1} \geq v_{i,j-1} + v_{i-1,j}$ , which is equivalent to the condition  $c_{i,j} \geq 0$ .

In the other direction, consider any two points  $(i, j)$  and  $(i', j')$  such that  $0 \leq i < i' \leq n$ ,  $0 \leq j < j' \leq n$ . Using Claim 2 we obtain

$$\begin{aligned} &v_{i',j'} - v_{i',j} - v_{i,j'} + v_{i,j} \\ &= \sum_{k=0}^{i'} \sum_{\ell=0}^{j'} c_{k,\ell} - \sum_{k=0}^{i'} \sum_{\ell=0}^j c_{k,\ell} - \sum_{k=0}^i \sum_{\ell=0}^{j'} c_{k,\ell} + \sum_{k=0}^i \sum_{\ell=0}^j c_{k,\ell} \\ &= \sum_{k=i+1}^{i'} \sum_{\ell=j+1}^{j'} c_{k,\ell} \end{aligned} \quad (4)$$



But  $C_V$  is non-negative and therefore  $v_{i',j'} - v_{i',j} - v_{i,j'} + v_{i,j} \geq 0$  as required. ■

It follows from Claim 3 that if we find some entry of  $C_V$  that is negative, then we have evidence that  $V$  is not an inverse Monge matrix. However, it is not necessarily true that if  $V$  is far from being an inverse Monge matrix, then  $C_V$  contains many negative entries. For example, suppose that  $C_V$  is 1 in all entries except the entry  $c_{n/2,n/2}$  which is  $-n^2$ . Then it can be verified that  $V$  is very far from being an inverse Monge matrix (this can be proved by showing that there are  $\Theta(n^2)$  disjoint quadruples  $v_{i,j}, v_{i',j'}, v_{i,j'}, v_{i',j}$  in  $V$ , such that from any such quadruple at least one value should be changed in order to transform  $V$  into an inverse Monge matrix). However, as our analysis will show, in such a case there are many sub-matrices in  $C_V$  whose sum of elements is negative. Thus our testing algorithms will sample certain sub-matrices of  $C_V$  and check that the sum of elements in each sub-matrix sampled is non-negative. We first observe that it is possible to check this efficiently.

**Claim 4** *Given access to  $V$  it is possible to check in time  $O(1)$  if the sum of elements in a given sub-matrix  $A$  of  $C_V$  is non-negative. In particular, if the lower-left entry of  $A$  is  $(i, j)$  and its upper-right entry is  $(i', j')$  then the sum of elements of  $A$  is  $v_{i',j'} - v_{i',j-1} - v_{i-1,j'} + v_{i-1,j-1}$ .*

**Proof:** Assume that  $A = (c_{k,\ell})_{k=i,\ell=j}^{k=i',\ell=j'}$  is a sub-matrix of  $C_V$ . Recall that for any  $q, p$ , we have  $v_{q,p} = \sum_{k=0}^q \sum_{\ell=0}^p c_{k,\ell}$ . Thus the sum of elements of  $A$  is:

$$\begin{aligned} \sum_{k=i}^{i'} \sum_{\ell=j}^{j'} c_{k,\ell} &= \sum_{k=0}^{i'} \sum_{\ell=j}^{j'} c_{k,\ell} - \sum_{k=0}^{i-1} \sum_{\ell=j}^{j'} c_{k,\ell} \\ &= \left( \sum_{k=0}^{i'} \sum_{\ell=0}^{j'} c_{k,\ell} - \sum_{k=0}^{i'} \sum_{\ell=0}^{j-1} c_{k,\ell} \right) - \left( \sum_{k=0}^{i-1} \sum_{\ell=0}^{j'} c_{k,\ell} - \sum_{k=0}^{i-1} \sum_{\ell=0}^{j-1} c_{k,\ell} \right) \\ &= (v_{i',j'} - v_{i',j-1}) - (v_{i-1,j'} - v_{i-1,j-1}). \end{aligned}$$

Therefore computing the sum of elements of any sub-matrix  $A$  of  $C_V$ , can be done by checking only 4 entries in the matrix  $V$ . ■

### 3.1 Filling Sub-matrices

An important building block for the analysis of our algorithms is a procedure for “filling in” a sub-matrix. That is, given constraints on the sum of elements in each row and column of a given sub-matrix, we are interested in assigning values to the entries of the sub-matrix so that these constraints are met.

Specifically, let  $a_1, \dots, a_s$  and  $b_1, \dots, b_t$  be non-negative real numbers such that  $\sum_{i=1}^s a_i \geq \sum_{j=1}^t b_j$ . Then it is possible to construct an  $s \times t$  non-negative real matrix  $T$ , such that the sum of elements in column  $j$  is exactly  $b_j$  and the sum of elements in row  $i$  is at most  $a_i$ . In the special case that  $\sum_{i=1}^s a_i = \sum_{j=1}^t b_j$ , the sum of elements in row  $i$  will equal  $a_i$ . In particular, this can be done by applying the following procedure, which is the same as the one applied to obtain an initial feasible solution for the linear-programming formulation of the transportation problem.

**Procedure 1** [Fill Matrix  $T = (t_{i,j})_{i,j=1}^{i=s,j=t}$ ]

Initialize  $\bar{a}_i = a_i$  for  $i = 1, \dots, s$  and  $\bar{b}_j = b_j$  for  $j = 1, \dots, t$ .

(In each of the following iterations,  $\bar{a}_i$  is an upper bound on what remains to be filled in row  $i$ ,

and  $\bar{b}_j$  is what remains to be filled in column  $j$ .)

for  $j = 1, \dots, t$ :

for  $i = 1, \dots, s$ :

Assign to entry  $(i, j)$  the value  $x = \min\{\bar{a}_i, \bar{b}_j\}$

Update  $\bar{a}_i = \bar{a}_i - x$ ,  $\bar{b}_j = \bar{b}_j - x$ .

**Claim 5** Procedure 1 fills the matrix  $T$  with non-negative values  $t_{i,j}$ , such that at the end of the procedure,  $\sum_{i=1}^s t_{i,j} = b_j$  for every  $j = 1, \dots, t$ , and  $\sum_{j=1}^t t_{i,j} \leq a_i$  for every  $i = 1, \dots, s$ . If initially  $\sum_{j=1}^t b_j = \sum_{i=1}^s a_i$  then  $\sum_{j=1}^t t_{i,j} = a_i$  for every  $i = 1, \dots, s$ .

**Proof:** Notice that initially  $\bar{a}_i = a_i \geq 0$  and  $\bar{b}_j = b_j \geq 0$ . Thus when we update  $\bar{a}_i = \bar{a}_i - x = \bar{a}_i - \min\{\bar{a}_i, \bar{b}_j\} \geq 0$  and similarly  $\bar{b}_j = \bar{b}_j - x = \bar{b}_j - \min\{\bar{a}_i, \bar{b}_j\} \geq 0$ . Therefore the  $\bar{a}_i$ 's and  $\bar{b}_j$ 's are always non-negative. Hence all values  $x$  filled in  $T$  are non-negative, since  $x = \min\{\bar{a}_i, \bar{b}_j\} \geq 0$ . Furthermore, after each such update the new sum over the  $\bar{a}_i$ 's equals the old sum over the  $\bar{a}_i$ 's minus  $x$  and a similar statement holds for the sum over the  $\bar{b}_j$ 's. Thus at all stages of the procedure,  $\sum_{i=1}^s \bar{a}_i \geq \sum_{j=1}^t \bar{b}_j$ , and if initially  $\sum_{i=1}^s a_i = \sum_{j=1}^t b_j$  then  $\sum_{i=1}^s \bar{a}_i = \sum_{j=1}^t \bar{b}_j$ .

We now show that the sum of elements in each column is as required. Observe that the procedure fills the columns one by one. Therefore when we start to fill column  $j$  we have  $\bar{b}_j = b_j$ . Since  $\sum_{i=1}^s \bar{a}_i \geq \sum_{j=1}^t \bar{b}_j$  at this stage, and all  $\bar{a}_i$ 's are non-negative, necessarily,  $\sum_{i=1}^s \bar{a}_i \geq \bar{b}_j = b_j$ . Let  $1 \leq k \leq s$  be the minimum integer such that  $\sum_{i=1}^k \bar{a}_i \geq b_j$ . Then by definition of the procedure, for every  $i < k$ , the entry  $(i, j)$  is filled with the value  $\bar{a}_i$ , and the entry  $(k+1, j)$  is filled with the value  $b_j - \sum_{i=1}^k \bar{a}_i$ . The total is hence  $b_j$  as required.

As for the rows, at all stages  $\bar{a}_i$  equals  $a_i$  minus the sum of all elements filled so far in row  $i$ . Therefore since  $\bar{a}_i \geq 0$ , then the sum of elements in row  $i$  is at most  $a_i$ . Furthermore, if initially  $\sum_{i=1}^s a_i = \sum_{j=1}^t b_j$ , then the sum of elements in row  $i$  will be exactly  $a_i$ . To show this note that at the end of the procedure,  $\sum_{j=1}^t \bar{b}_j = 0$ , since each  $\bar{b}_j$  equals  $b_j$  minus the sum of all elements in column  $j$ , and we have shown that the sum of elements in column  $j$  is  $b_j$ . But  $\sum_{i=1}^s \bar{a}_i = \sum_{j=1}^t \bar{b}_j$ , and therefore also  $\sum_{i=1}^s \bar{a}_i = 0$  at the end. Since  $\bar{a}_i \geq 0$ , this means that  $\bar{a}_i = 0$ . Hence the sum of elements in row  $i$  must be  $a_i$ . ■

## 4 A Testing Algorithm for Inverse Monge Matrices

We first present a simple algorithm for testing if a matrix  $V$  is an inverse Monge Matrix, whose running time is  $O(n/\epsilon)$ . In the next section we show a significantly faster algorithm that is based on the ideas presented here. We may assume without loss of generality that  $n$  is a power of 2. This is true since our algorithms probe the coefficients matrix  $C_V$ , and we may simply “pad” it by 0's to obtain rows and columns that have lengths which are powers of 2 and run the algorithm with  $\epsilon \leftarrow \epsilon/4$ . We shall need the following two definitions for both algorithms.

**Definition 4 (Sub-Rows, Sub-Columns and Sub-Matrices)** A sub-row in an  $n \times n$  matrix is a consecutive sequence of entries that belong to the same row. The sub-row  $((i, j), (i, j+1), \dots, (i, j+t-1))$  is denoted by  $[ ]_{i,j}^{1,t}$ . A sub-column is defined analogously, and is denoted by  $[ ]_{i,j}^{s,1} = ((i, j), (i+1, j), \dots, (i+s-1, j))$ . More generally, an  $s \times t$  sub-matrix whose bottom-left entry is  $(i, j)$  is denoted  $[ ]_{i,j}^{s,t}$ .

**Definition 5 (Legal Sub-Matrices)** *A sub-row in an  $n \times n$  matrix is a legal sub-row if it can result from bisecting the row of length  $n$  that contains it in a recursive manner. That is, a complete (length  $n$ ) row is legal, and if  $[ ]_{i,j}^{1,t}$  is legal, then so are  $[ ]_{i,j}^{1,t/2}$  and  $[ ]_{i,j+t/2}^{1,t/2}$ . A legal sub-column is defined analogously. A sub-matrix is legal if both its rows and its columns are legal.*

Note that the legality of a sub-row  $[ ]_{i,j}^{1,t}$  is independent of the actual row  $i$  it belongs to, but rather it depends on its starting position  $j$  and ending position  $j + t - 1$  within its row. An analogous statement holds for legal sub-columns.

Although a sub-matrix is just a collection of positions (entries) in an  $n \times n$  matrix, we talk throughout the paper about sums of elements in certain sub-matrices  $A$  of  $C_V$ . In this we mean the sum of elements of  $C_V$  determined by the set of positions in  $A$ .

**Definition 6 (Good sub-matrix)** *We say that a sub-matrix  $A$  of  $C_V$  is good if the sum of elements in each row and each column of  $A$  is non-negative.*

**Definition 7 (Good Point)** *We say that point  $(i, j)$  is good if all legal square sub-matrices  $A$  of  $C_V$  which contain  $(i, j)$  are good.*

**Algorithm 2 [Test Monge I]**

1. Choose  $8/\epsilon$  points in the matrix  $C_V$  and check that they are good.
2. If all points are good then accept, otherwise reject.

By Claim 4, it is possible to check in constant time that the sum of elements in a sub-row (sub-column) of  $C_V$  is non-negative. Therefore, it is possible to test that an  $s \times s$  square sub-matrix  $A$  of  $C_V$  is good in time  $\Theta(s)$ . Notice that every point in an  $n \times n$  matrix is contained in  $\log n$  square sub-matrices. Hence the time required to check whether a point is good is  $O(n) + O(n/2) + \dots + O(n/2^i) + \dots + O(1) = O(n)$ , and the complexity of the algorithm is  $O(n/\epsilon)$ .

**Theorem 2** *If  $V$  is an inverse Monge matrix then it is always accepted, and if  $V$  is  $\epsilon$ -far from being an inverse Monge matrix, then the algorithm rejects with probability at least  $2/3$ .*

**Proof:** The first part of the theorem follows directly from Claim 3. In order to prove the second part of the theorem, we show that if  $V$  is  $\epsilon$ -far from being inverse Monge, then  $C_V$  contains more than  $(\epsilon/4)n^2$  bad points. The second part of the theorem directly follows because the probability in such a case that no bad point is selected by the algorithm, is at most  $(1 - \epsilon/4)^{(8/\epsilon)} < e^{-2} < 1/3$ .

Assume contrary to the claim that  $C_V$  contains at most  $(\epsilon/4)n^2$  bad points. We shall show that by modifying at most  $\epsilon n^2$  entries in  $V$  we obtain an inverse Monge matrix (in contradiction to our assumption concerning  $V$ ). Let us look at the set of bad points in  $C_V$ , and for each such bad point look at the largest bad square sub-matrix in  $C_V$  which contains this bad point. By our assumption on the number of bad points, it must be the case that the area of all these maximal bad sub-matrices is at most  $(\epsilon/4)n^2$ , because all the points in a bad sub-matrix are bad.

For each maximal bad (legal square) sub-matrix  $B$  of  $C_V$  we will look at the smallest good (legal square) sub-matrix  $A$  which contains  $B$ . First observe that such a good sub-matrix must exist. Indeed, since  $B$  is maximal, if it is of size  $s \times s$  where  $s < n$ , then the legal square sub-matrix of size  $2s \times 2s$  that contains it must be good. But if  $s = n$ , then  $B = C_V$  implying that all  $n^2$  points in  $C_V$  are bad, contradicting our assumption on the number of bad points. Next observe that for every two maximal sub-matrices  $B$  and  $B'$ , the corresponding good sub-matrices  $A$  and  $A'$

that contain them are either the same sub-matrix, or are totally disjoint. Finally, the sum of areas of all these good sub-matrices is at most  $4 \cdot (\epsilon/4)n^2 = \epsilon n^2$ .

We now correct each such good sub-matrix  $A$  so that it contains only non-negative elements, and the sum of elements in each row and column of  $A$  remains as it was. This can be done by applying Procedure 1 to  $A$  as described in Section 3.1.

Note that after correcting all these good sub-matrices of  $C_V$ , the new matrix  $C_V$  is non-negative, and thus the corresponding new matrix  $V$  must be an inverse Monge matrix. We must show however, that at most  $\epsilon n^2$  values were changed in  $V$  following the changes to  $C_V$ . Notice that we made sure that the sum of elements in each row and column of each corrected sub-matrix  $A$  remains as it was. Therefore the values of all points  $v_{k,\ell}$  in  $V$  that are outside  $A$  are not affected by the change to  $A$ , since by Claim 2 we have that  $v_{k,\ell} = \sum_{i=0}^k \sum_{j=0}^{\ell} c_{i,j}$ . ■

## 5 A Faster Algorithm for Inverse Monge Matrices

Though the above algorithm has running time sub-linear in the size of the matrix, which is  $n^2$ , we would further like to improve its dependence on  $n$ . We next suggest a variant of the algorithm whose running time is  $O(\log^2 n/\epsilon)$  and explain what needs to be proved in order to argue its correctness. We first redefine the concepts of good sub-matrices and good points.

**Definition 8 (Good sub-matrix)** *A (legal) sub-matrix  $T$  of  $C_V$  is good if the sum of all its elements is non-negative. Otherwise,  $T$  is bad.*

**Definition 9 (Good Point)** *We say that a point is good if every legal sub-matrix of  $C_V$  that contains it is good. Otherwise the point is bad.*

For the sake of the presentation, we shall assume that every row and every column in  $C_V$  (that is, every sub-row and sub-column of length  $n$ ) have a non-negative sum. In Subsection 5.2 we explain how to remove this assumption. Note that this assumption implies that every  $s \times n$  sub-matrix is good, and similarly for every  $n \times s$  sub-matrix (but of course it has no implications on smaller sub-matrices).

### Algorithm 3 [Test Monge II]

1. Uniformly select  $8/\epsilon$  points in the matrix  $C_V$  and check that they are good.
2. If all points are good then accept, otherwise reject.

Note that by Definition 5, each point in an  $n \times n$  matrix is contained in  $O(\log^2 n)$  legal sub-matrices. Thus by Claim 4, checking that a point is good takes time  $O(\log^2 n)$ . Therefore the running time of the algorithm is  $O((\log^2 n)/\epsilon)$ .

**Theorem 3** *If  $V$  is an inverse Monge matrix then it is always accepted, and if  $V$  is  $\epsilon$ -far from being an inverse Monge matrix, then the algorithm rejects with probability at least  $2/3$ .*

### 5.1 Outline of the Proof of Theorem 3

If  $V$  is an inverse Monge matrix then by Claim 3 all elements in  $C_V$  are non-negative, and so the algorithm always accepts. Suppose  $V$  is  $\epsilon$ -far from being inverse Monge. We claim that in such a case  $C_V$  must contain more than  $(\epsilon/4)n^2$  bad points, causing the algorithm to reject with

probability at least  $1 - (1 - \epsilon/4)^{(8/\epsilon)} > 1 - e^{-2} > 2/3$ . Assume contrary to the claim that  $C_V$  contains at most  $(\epsilon/4)n^2$  bad points. Our goal from this point on is to show that in such a case  $V$  is  $\epsilon$ -close to being an inverse Monge matrix.

Consider the union of all bad legal sub-matrices of  $C_V$ . Since within each bad legal sub-matrix, all points are bad, then the area occupied by this union is at most  $(\epsilon/4)n^2$ .

**Definition 10 (Maximal bad legal sub-matrix)** *A bad legal sub-matrix  $T$  of  $C_V$  is a maximal bad legal sub-matrix of  $C_V$  if it is not contained in any larger bad legal sub-matrix of  $C_V$ .*

Now consider all such maximal bad legal sub-matrices of  $C_V$ . For each such sub-matrix  $B$  let us take the legal sub-matrix that contains it and has twice the number of rows and twice the number of columns. Then by the maximality of  $B$  (and our assumption that all full rows and columns have a non-negative sum), the resulting sub-matrix is good. We now take the union of all these good legal sub-matrices, and get a total area of size at most  $\epsilon n^2$ . Denote the union of all these sub-matrices by  $R$ . See for example Figure 1.




Figure 1: An example of the structure of a subset  $R$  (outlined by a bold line). The bad legal sub-matrices determining  $R$  are the gray sub-matrices. Each is contained inside a good legal sub-matrix that has twice the number of rows and twice the number of columns (marked by dashed rectangles). Observe that maximal bad-legal sub-matrices may overlap.

**Definition 11 (Maximal (legal) sub-row/column)** *Given a subset  $R$  of entries in an  $n \times n$  matrix, a sub-row  $T$  is a maximal (legal) sub-row with respect to  $R$  if  $T$  is contained in  $R$  and there is no larger (legal) sub-row  $T'$  such that  $T \subset T' \subseteq R$ . A maximal (legal) sub-column with respect to  $R$  is defined analogously.*

For sake of succinctness, whenever it is clear what  $R$  is, we shall just say maximal (legal) sub-row and drop the suffix, “with respect to  $R$ ”. Note that a maximal sub-row is simply a maximal

consecutive sequence of entries in  $R$  that belong to the same row, while a maximal legal sub-row is a more constrained notion. In particular, a maximal sub-row may be a concatenation of several maximal legal sub-rows.

We would like to show that it is possible to change the at most  $\epsilon n^2$  entries of  $C_V$  within  $R$  to non-negative values so that the following property holds:

**Property 1 (Sum Property for  $R$ )** *For every point  $(i, j)$  outside of  $R$ , the sum of the elements in the modified entries  $(i', j')$  within  $R$  such that  $i' \leq i$  and  $j' \leq j$  is as it was in the original matrix  $C_V$ .*

Let  $\tilde{C}_V$  be the matrix obtained from  $C_V$  by modifying  $R$  so that Property 1 holds, and let  $\tilde{V}$  be the matrix which corresponds to  $\tilde{C}_V$ . Then it follows from Claim 2 that  $\tilde{V}$  is at most  $\epsilon$ -far from the original matrix  $V$ .

## 5.2 Dealing with Rows/Columns Having a Negative Sum

Before we continue with showing how to obtain Property 1, we explain shortly how to remove the assumption that all rows and columns in  $C_V$  have a non-negative sum. If  $\epsilon \leq 4/n$  then we may directly check in time  $O(1/\epsilon)$  that in fact all rows and columns of the matrix  $C_V$  have a non-negative sum (using Claim 4), and reject if some row or column has a negative sum. Hence in this case our assumption is valid. Thus assume that  $\epsilon > 4/n$ .

First we slightly modify Algorithm 3 so that it uniformly selects  $16/\epsilon$  points in  $C_V$  (instead of  $8/\epsilon$ ). In such a case, if  $C_V$  contains more than  $(\epsilon/8)n^2$  bad points then the algorithm rejects with probability at least  $2/3$ . We thus assume that  $C_V$  contains at most  $(\epsilon/8)n^2$  bad points and strive to show that in such a case  $V$  is  $\epsilon$ -close to being an inverse Monge matrix. Since we do not assume that every row and column in  $C_V$  has a non-negative sum, we first modify  $C_V$  so that it has this property.

Consider each row  $i$  in  $C_V$  whose sum of entries is negative. Suppose that we modify the last entry in the row,  $c_{i,n}$ , so that the new sum of all entries is 0. Similarly, we modify the last entry  $c_{n,j}$  in each column  $j$  that has a negative sum. Let  $\bar{C}_V$  be the resulting coefficients matrix, and let  $\bar{V}$  be the corresponding value matrix. Then all rows and columns in  $\bar{C}_V$  have a non-negative sum, and by Claim (2)  $\bar{V}$  and  $V$  differ on at most  $2n - 1 < (\epsilon/2)n^2$  entries (at most all entries in the last column and last row).

Now we may define the region  $R$  as we did in the previous subsection. Note that in this case the area of the region  $R$  is at most  $(\epsilon/2)n^2$ . We can therefore continue in proving that it is possible to modify only the entries within  $R$  so that they are all positive and Property 1 holds. This will imply that the total number of entries that should be modified (first to obtain non-negative rows and columns, and then to “fix”  $R$ ) is at most  $\epsilon n^2$ , as desired.

## 5.3 Fixing $R$

Let  $R$  be the subset of entries in the matrix  $C_V$  that consists of a union of good legal sub-matrices. In the following discussion, when we talk about elements in sub-matrices of  $R$  we mean the elements in  $C_V$  determined by the corresponding set of positions in  $R$ .

**Lemma 6** *The sum of elements in every maximal legal sub-row and every maximal legal sub-column in  $R$  is non-negative.*

**Proof:** Assume, contrary to the claim, that  $R$  contains some maximal legal sub-row  $L = [ ]_{i,j}^{1,t}$  whose sum of elements is negative. Let  $T$  be the maximal bad legal sub-matrix in  $C_V$  that contains  $L$ . By the maximality of  $L$ , necessarily  $T = [ ]_{i',j}^{s,t}$  for some  $i' \leq i$  and  $s \geq 1$ . That is, the rows of  $T$  (one of which is  $L$ ) are of length  $t$ . By the construction of  $R$ ,  $R$  must contain a good legal sub-matrix  $T'$  that contains  $T$  and is twice as large in each dimension. But this contradicts the maximality of  $L$ . ■

### 5.3.1 Maximal Blocks

We will partition  $R$  into disjoint blocks (sub-matrices) and fill each block separately with non-negative values, so that the sum property for  $R$  is maintained (see Property 1). We define blocks as follows.

**Definition 12 (Maximal Block)** *A maximal block  $B = [ ]_{i,j}^{s,t}$  in  $R$  is a sub-matrix contained in  $R$  which has the following property: It consists of a maximal consecutive sequence of maximal legal sub-columns of the same height. The maximality of each sub-column is as in Definition 11. That is, for every  $j \leq r \leq j + t - 1$ , the column  $[ ]_{i,r}^{s,1}$  is a maximal legal sub-column (with respect to  $R$ ).*

The maximality of the sequence of sub-columns in a block means that we cannot extend the sequence of columns neither to the left nor to the right. That is, neither  $[ ]_{i,j-1}^{s,1}$  nor  $[ ]_{i,j+t}^{s,1}$  are maximal legal sub-columns in  $R$ . (Specifically, each is either not fully contained in  $R$  or  $R$  contains a larger legal sub-column that contains it.)

We shall sometimes refer to maximal blocks simply as blocks. Observe that by this definition,  $R$  is indeed partitioned in a unique way into maximal disjoint blocks. See Figure 2 for an example of  $R$  and its partition into maximal blocks.

**Definition 13 (Size of a Maximal Block)** *Let  $B$  be a maximal block. The size of  $B$  is the height of the columns in  $B$  (equivalently, the number of rows in  $B$ ).*

### 5.3.2 Bounded Sub-Matrices

As we have shown in Lemma 6, the sum of elements in every maximal legal sub-column in  $R$  is non-negative. It directly follows that every maximal block has a non-negative sum. We would like to characterize other sub-matrices of  $R$  whose sum is necessarily non-negative.

**Definition 14** *For a given sub-matrix  $T$ , we denote the sum of the elements in  $T$  by  $sum(T)$ .*

**Lemma 7** *Consider any two maximal blocks  $B = [ ]_{i,j}^{s,t}$  and  $B' = [ ]_{i,j'}^{s,t'}$  where  $j' > j + t$ . That is, both blocks have the same size  $s$  and both start at row  $i$  and end at row  $i + s - 1$ . Consider the sub-matrix  $T = [ ]_{i,j+t}^{s,j'-(j+t)}$  "between them". Suppose that  $T \subset R$ . Then  $sum(T) \geq 0$ .*

**Proof:** Consider first the case that  $T$  is a legal sub-matrix. Assume, contrary to the claim, that  $sum(T) < 0$ . That is,  $T$  is a bad legal sub-matrix. Let  $T'$  be the maximal bad legal sub-matrix containing  $T$ . By construction of  $R$ ,  $R$  should contain a good legal sub-matrix  $T''$  that contains  $T'$  and has twice the number of rows and twice the number of columns. But this would contradict the maximality of the sub-columns of  $B$  or of  $B'$ . To see why this is true, assume without loss of generality that for any legal sub-column  $[ ]_{i,r}^{s,1}$ , the legal column that is twice its size is  $[ ]_{i,r}^{2s,1}$ . Then  $T''$  must contain either the sub-column  $[ ]_{i,j'}^{2s,1}$  or the sub-column  $[ ]_{i,j+t-1}^{2s,1}$  (depending on the identity of the legal sub-rows that are twice the size of the rows of  $T'$ ). In the first case we would




Figure 2: An example of the partition of  $R$  into maximal blocks (numbered  $B_1$ – $B_9$ ). Note that the ratio between the sizes of any two blocks is always a power of 2. Furthermore, the blocks are “oriented” in the following sense. Suppose a block  $B$  has size  $s$  and a block  $B'$  has size  $s' \leq s$  and some of their sub-rows belong to the same row of the matrix (e.g.,  $B_4$  and  $B_2$ , or  $B_9$  and  $B_6$ ). Then the smaller block  $B'$  must be aligned either with the first or second half of  $B$ , or with one of the quarters of  $B$ , or with one of its eighth's, etc.

get a contradiction to the fact that  $B$  is a maximal block, and in the second case we would get a contradiction of the fact that  $B'$  is a maximal block.

Suppose that  $T$  is not a legal sub-matrix. Observe that its columns are necessarily legal sub-columns (given that the columns of  $B$  and  $B'$  are legal). Hence, only its rows are not legal sub-rows. Therefore,  $T$  can be partitioned into sub-matrices  $T_1, \dots, T_k$ , such that each is of height  $s$ , and is a maximal legal sub-matrix *with respect to*  $T$ . We claim that for every  $T_\ell$ ,  $\text{sum}(T_\ell) \geq 0$ . Consider any fixed  $T_\ell$ . By its maximality with respect to  $T$ , we know that the legal sub-rows that contain the rows of  $T_\ell$  and are twice their size, are not in  $T$  and hence must intersect either  $B$  or  $B'$ . Thus, if we let  $T'_\ell$  be the maximal bad legal sub-matrix with respect to  $R$  that contains  $T_\ell$ , and we let  $T''_\ell$  be the good legal sub-matrix that contains  $T'_\ell$  and is twice its size, then we reach a contradiction with the maximality of either  $B$  or  $B'$ . ■

More generally we can obtain:

**Lemma 8** *Consider any two maximal blocks  $B = [ ]_{i,j}^{s,t}$  and  $B' = [ ]_{i',j'}^{s',t'}$  where  $i \leq i' \leq i + s - 1$ ,  $i' + s' \leq i + s$ , and  $j' \geq j + t$ , or  $j' + t' \leq j$ . That is,  $B$  has size  $s$  and  $B'$  has size  $s' \leq s$ , and  $B'$  starts at row  $i' \geq i$  and ends at row  $i + s' - 1 \leq i + s - 1$ . Consider the sub-matrix  $T$  of height  $s$  “between them”, so that  $T = [ ]_{i,j+t}^{s,j'-(j+t)}$  or  $T = [ ]_{i,j'+t}^{s,j-(j'+t)}$ . Suppose that  $T \subset R$ . Then  $\text{sum}(T) \geq 0$ .*

**Proof:** If  $s' = s$  then we have the case treated in Lemma 7. Hence assume  $s' < s$ . The proof of this case follows exactly the same outline as the proof of Lemma 7. The only point that needs to be noticed is the following. Recall that both the columns of  $B$  and of  $B'$  are legal sub-columns, and that  $i \leq i' \leq i + s - 1$ ,  $i' + s' \leq i + s$ . Also recall that the ratio between the sizes of any two






Figure 3: Illustrations for Lemmas 7 and 8. The first figure illustrates the case, in the proof of Lemma 7, where  $T$  is a legal sub-matrix (for simplicity, we assume  $T' = T$ ). The second figure illustrates the second case in the proof when  $T$  is a union of legal sub-matrices (all having the height of  $B$ ). The third figure illustrates the case Lemma 8 deals with when  $B$  and  $B'$  are of a different size.

blocks is always a power of 2. Furthermore, the blocks are “oriented” in the sense that  $B'$  must be aligned either with the first or second half of  $B$ , or with one of the quarters of  $B$ , or with one of its eighths, etc. Hence, for every column  $[ ]_{i',j''}^{s',1}$  of  $B'$ , the legal column that contains it and is twice its size, is contained within  $[ ]_{i,j''}^{s,1}$ . Hence, if  $T''$  is an extension of  $T$  to the direction of  $B'$ , so that it contains either  $[ ]_{i,j'}^{s,1}$  or  $[ ]_{i,j'+t'-1}^{s,1}$ , then we necessarily get a contradiction of the maximality of the sub-columns of  $B'$ . ■

See Figure 3 for illustrations of Lemmas 7 and 8.

**Definition 15 (Covers)** We say that a collection  $A$  of sub-rows covers a given block  $B$  with respect to  $R$ , if  $B \subset A \subset R$  and the number of rows in  $A$  equals the size of  $B$ . We say that  $A$  is a maximal row-cover with respect to  $R$  if  $A$  consists of maximal sub-rows with respect to  $R$ .

**Definition 16 (Borders)** We say that a sub-matrix  $A = [ ]_{i,j}^{s,t}$  in  $R$ , borders a maximal block  $B = [ ]_{i',j'}^{s',t'}$  if  $i \leq i' \leq i + s - 1$ ,  $i' + s' \leq i + s$ , and either  $j' = j + t$  (so that  $A$  borders  $B$  from the left), or  $j' + t' = j$  (so that  $A$  borders  $B$  from then right).

By Lemmas 7 and 8, and using the above terminology, we get the following corollary whose proof is illustrated in Figure 4.




Figure 4: An illustration for Corollary 9. Here  $A$  covers the blocks  $B_1$ ,  $B_2$  and  $B_3$ , and borders the blocks  $D_1$  and  $D_2$ . The sub-matrices  $T_0$ – $T_4$  are parts of larger blocks (that extend above and/or below  $A$ ).

**Corollary 9** *Let  $A$  be a sub-matrix of  $R$  which covers a given block  $B$ . If on each of its sides  $A$  either borders a block smaller than  $B$  or its border coincides with the border of  $R$ , then  $sum(A) \geq sum(B)$ .*

**Proof:** Let  $B_1, \dots, B_k$  be the set of maximal blocks that are covered by  $A$  (where  $B = B_i$  for some  $1 \leq i \leq k$ ). Note that by definition of maximal blocks, they are all of the same size (which is the height of  $A$ ). Let  $D_1$  and  $D_2$  be the smaller blocks that border  $A$  on the left side and the right side respectively. (If there is no such block on one of the sides, then we think of the corresponding  $D_i$  as having size 0). Let  $T_0, \dots, T_k$  be the sub-matrices between these blocks. That is,  $T_0$  is between  $D_1$  and  $B_1$ ,  $T_k$  is between  $B_k$  and  $D_2$ , and for  $1 \leq i \leq k - 1$ ,  $T_i$  is between  $B_i$  and  $B_{i+1}$ . Then, by Lemmas 7 and 8

$$sum(A) = \sum_{i=1}^k sum(B_i) + \sum_{i=0}^k sum(T_i) \geq sum(B). \quad (5)$$

■

### 5.3.3 The Procedure for Refilling $R$

We now describe the procedure that refills the entries of  $R$  with non-negative values. Recall that  $R$  is a disjoint union of maximal blocks. Hence if we remove a maximal block from  $R$ , then the maximal blocks of the remaining structure, are simply the remaining maximal blocks of  $R$ . The procedure described below will remove the blocks of  $R$  one by one, in order of increasing size, and refill each block separately using Procedure 1. After removing each block, the sum of the elements in each remaining column in  $R$  remains the same, however the row sums must be updated. Procedure 1 is used here as well.

#### Procedure 2 [Refill $R$ ]

1. We assign with each maximal sub-row  $L$  in  $R$  a designated sum of elements for that row, which is denoted by  $\overline{sum}(L)$ , and initially set to be  $\overline{sum}(L) = sum(L)$ .
2. Let  $m$  be the number of maximal blocks in  $R$ , and let  $R_1 = R$ .
3. for  $p = 1, \dots, m$ :

- (a) Let  $B_p$  be a maximal block in  $R_p$  whose size is minimum among all maximal blocks of  $R_p$ , and assume that  $B_p$  is an  $s \times t$  sub-matrix. Let  $A_p$  be a maximal row-cover of  $B_p$  with respect to  $R_p$ . For  $1 \leq \ell \leq s$ , let  $L_\ell$  denote the sub-row of  $A_p$  that covers the  $\ell$ 'th sub-row of  $B_p$ .
- (b) Refill  $B_p$  by applying Procedure 1 (see Section 3.1), where the sum filled in the  $k$ 'th sub-column of  $B_p$ ,  $1 \leq k \leq t$ , should be the original sum of this sub-column in  $C_V$ , and the sum filled in the  $\ell$ 'th sub-row of  $B_p$ ,  $1 \leq \ell \leq s$ , is at most  $\overline{\text{sum}}(L_\ell)$ .  
For each  $1 \leq \ell \leq s$ , let  $x_\ell$  denote the sum of elements filled by Procedure 1 in the  $\ell$ 'th sub-row of  $B_p$ .

- (c) Let  $R_{p+1} = R_p \setminus B_p$  and assign designated sums to the rows of  $R_{p+1}$  that have been either shortened or broken into two parts by the removal of  $B_p$  from  $R_p$ . This is done as follows:

The set  $A_p \setminus B_p$  is the union of two non-consecutive sub-matrices,  $A'$  and  $A''$ , so that  $A'$  borders  $B_p$  from the left and  $A''$  borders  $B_p$  from the right. Let  $L'_\ell$  and  $L''_\ell$  be the sub-rows in  $A'$  and  $A''$  respectively that are contained in sub-row  $L_\ell$  of  $A_p$ . We assign to  $L'_\ell$  and  $L''_\ell$  non-negative designated sums,  $\overline{\text{sum}}(L'_\ell)$  and  $\overline{\text{sum}}(L''_\ell)$ , that satisfy the following:

$$\overline{\text{sum}}(L'_\ell) + \overline{\text{sum}}(L''_\ell) = \overline{\text{sum}}(L_\ell) - x_\ell,$$

and furthermore,

$$\sum_{\text{row } L \in A'} \overline{\text{sum}}(L) = \text{sum}(A'), \quad \sum_{\text{row } L \in A''} \overline{\text{sum}}(L) = \text{sum}(A'').$$

This is done by applying Procedure 1 to a  $2 \times s$  matrix whose sums of columns are  $\text{sum}(A')$  and  $\text{sum}(A'')$  and sums of rows are  $\overline{\text{sum}}(L_\ell) - x_\ell$ , where  $1 \leq \ell \leq s$ .

(Note that one or both of  $A'$  and  $A''$  may not exist. This can happen if  $B_p$  bordered  $A_p \setminus B_p$  on one side and its boundary coincided with  $R_p$ , or if  $A_p = B_p$ . In this case, if, for example,  $A'$  does not exist then we view it as a sub-matrix of size 0 where  $\text{sum}(A') = 0$ .)

## 5.4 Proving that Procedure 2 Works

Recall that for each  $1 \leq p \leq m$ ,  $R_p$  is what remains of  $R$  at the start of the  $p$ 'th iteration of Procedure 2. In particular,  $R_1 = R$ . We would first like to show that the procedure does not “get stuck”. That is, for each iteration  $p$ , Procedure 1 can be applied to the block  $B_p$  selected in this iteration, and the updating of the designated sum for the rows that have been shortened by the removal of  $B_p$  can be performed. Note that since the blocks are selected according to increasing size, then in each iteration the maximal row cover  $A_p$  of  $B_p$  must actually be a sub-matrix.

### 5.4.1 Proving that Procedure 2 Does not Get Stuck

For every  $1 \leq p \leq m$ , let  $s_p$  be the minimum size of the maximal blocks of  $R_p$ , where  $s_0 = 1$ . Observe that whenever  $s_p$  increases, it does so by a factor of  $2^k$  for some  $k$ . This is true because the columns of maximal blocks are legal sub-columns.

**Lemma 10** *For every  $1 \leq p \leq m$ , Procedure 1 can be applied to the block  $B_p$  selected in  $R_p$ , and the updating process of the designated sum of rows can be applied. Moreover, if  $A$  is a sub-matrix of  $R_p$  with height of at least  $s_{p-1}$ , whose columns are legal sub-columns and whose rows are maximal rows with respect to  $R_p$ , then  $\sum_{\text{row } L \in A} \overline{\text{sum}}(L) = \text{sum}(A)$ .*

**Proof:** Let  $B_p$  be the block selected in iteration  $p$ , where  $B_p$  is an  $s \times t$  sub-matrix, and let  $A_p$  be the maximal row-cover of  $B_p$  with respect to  $R_p$ . As noted in Subsection 3.1, all that is required for Procedure 1 to work is:

- (1) For every column  $K$  in  $B_p$ ,  $sum(K) \geq 0$ .
- (2) For every row  $L$  in  $A_p$ ,  $\overline{sum}(L) \geq 0$ .
- (3)  $\sum_{\text{row } L \in A_p} \overline{sum}(L) \geq \sum_{\text{column } K \in B_p} sum(K)$ .

In order for the updating process to succeed in Step 3c of Procedure 2, we must have that:

- (4) For each  $1 \leq \ell \leq s$ , let  $x_\ell$  be the sum of elements filled in the  $\ell$ 'th sub-row of  $B_p$ , and let  $L_\ell$  be the sub-row of  $A_p$  that covers this sub-row of  $B_p$ . Then,  $\overline{sum}(L_\ell) - x_\ell \geq 0$ .
- (5) If  $A_p \setminus B_p$  consists of the two sub-matrices  $A'$  and  $A''$  (between which resided  $B$ ), then  $sum(A') \geq 0$ ,  $sum(A'') \geq 0$ , and

$$\sum_{\text{row } L_\ell \in A_p} (\overline{sum}(L_\ell) - x_\ell) = sum(A') + sum(A'').$$

By Lemma 6, Item (1) holds at the start of every iteration. In order to prove the other items for every  $p$ , we first extend and generalize Item (2):

- (2') Let  $A$  be any sub-matrix in  $R_p$  having height at least  $s_{p-1}$  whose columns are legal sub-columns and whose rows are maximal rows with respect to  $R_p$ . Then for every row  $L$  of  $A$  we have  $\overline{sum}(L) \geq 0$ , and  $\sum_{\text{row } L \in A} \overline{sum}(L) = sum(A)$ .

Observe that if Item (2') holds at the start of iteration  $p$ , then in particular it holds for  $A_p$ . Hence by Corollary 9

$$\sum_{\text{row } L \in A_p} \overline{sum}(L) = sum(A_p) \geq sum(B_p) \tag{6}$$

and so Item (3) holds as well.

Furthermore, if Items (1)–(3) hold at the start of iteration  $p$ , then Procedure 1 can be applied successfully. Thus Item (4) necessarily holds by definition of Procedure 1. The first part of Item (5), concerning the non-negativity of  $A'$  and  $A''$ , follows from Lemmas 7 and 8 (similarly to the way Corollary 9 follows from these two lemmas). The second part of Item (5) follows from Item (2') holding for  $A_p$  and the fact that  $\sum_{\ell=1}^s x_\ell = sum(B_p)$  (since Procedure 1 completed successfully). Hence,

$$\sum_{\text{row } L_\ell \in A_p} (\overline{sum}(L_\ell) - x_\ell) = sum(A_p) - sum(B_p) = sum(A') + sum(A''). \tag{7}$$

as required.

Hence, it remains to prove that Item (2') holds at the start of every iteration  $p$ . We do so by induction on  $p$ . Consider the base case,  $p = 1$ , so that  $R_p = R_1 = R$ . By the initialization of Procedure 2, for every maximal sub-row  $L$  of  $R$ ,  $\overline{sum}(L) = sum(L)$ . By Lemma 6 (applied to the maximal legal sub-rows which partition  $L$ ), we know that  $\overline{sum}(L) \geq 0$ . Furthermore, every sub-matrix  $A$  of  $R$  having height of at least  $s_{p-1} = s_0 = 1$  is the union of maximal sub-rows of  $R$ , and so

$$\sum_{\text{row } L \in A} \overline{sum}(L) = \sum_{\text{row } L \in A} sum(L) = sum(A) \tag{8}$$

as required.

Assume the induction claim holds for  $p - 1$ , we prove it for  $p$ . Consider any sub-matrix  $A$  of height of at least  $s_{p-1}$ , whose columns are legal sub-columns and whose rows are maximal sub-rows with respect to  $R_p$ . If  $A$  also consisted of maximal sub-rows with respect to  $R_{p-1}$ , then we are done by the induction hypothesis.

Otherwise, the block  $B_{p-1}$  of size  $s_{p-1}$  that was removed from  $R_{p-1}$ , bordered  $A$  on one of its sides. Let  $A^1, \dots, A^q$  be the disjoint sub-matrices of height  $s_{p-1}$  such that  $A = \cup_{h=1}^q A^h$  (that is,  $A^1, \dots, A^q$  are located one on top of the other). In this case, all but at most one of these sub-matrices, say  $A^q$ , consisted of maximal sub-rows with respect to  $R_{p-1}$ , and  $B_{p-1}$  bordered  $A^q$ .

For each of the sub-matrices  $A^1, \dots, A^{q-1}$  we can apply the induction hypothesis (Item (2')). We get that for each such  $A^h$ : (a) For every row  $L$  in  $A^h$ ,  $\overline{sum}(L) \geq 0$ ; and (b)  $\sum_{\text{row } L \in A^h} \overline{sum}(L) = sum(A^h)$ .

As for  $A^q$ , assume without loss of generality that  $B_{p-1}$  bordered  $A^q$  from its right. Let  $A'$  be the sub-matrix that bordered  $B_{p-1}$  on its left ( $A'$  may be empty). This means that  $A_{p-1}$  is of the form  $A_{p-1} = A^q \cup B_{p-1} \cup A'$  (see Figure 5). But then, by definition of the updating rule and since it succeeded by the induction hypothesis (Items (4) and (5)), we have that for every row  $L$  in  $A^q$ ,  $\overline{sum}(L) \geq 0$  and  $\sum_{\text{row } L \in A^q} \overline{sum}(L) = sum(A^q)$ .

It follows that for every row  $L$  in  $A$  we have  $\overline{sum}(L) \geq 0$  and

$$\sum_{\text{row } L \in A} \overline{sum}(L) = \sum_{h=1}^q \sum_{\text{row } L \in A^h} \overline{sum}(L) = \sum_{h=1}^q sum(A^h) = sum(A) \quad (9)$$

and the induction step is proven. ■




Figure 5: An illustration for the induction step in the proof of Lemma 10 (where  $q = 4$ ).

#### 5.4.2 Proving that Procedure 2 is Correct

Let  $\tilde{C}_V = \{\tilde{c}_{i,j}\}$  be the matrix resulting from the application of Procedure 2 to the matrix  $C_V = \{c_{i,j}\}$ . For any sub-matrix  $T$  of  $C_V$  (and in particular of  $R$ ), we let  $\widetilde{sum}(T)$  denote the sum of elements of  $T$  in  $\tilde{C}_V$ . By definition of the procedure,  $\widetilde{sum}(K) = sum(K)$  for every maximal legal sub-column  $K$  of  $R$ . Hence this holds also for every maximal sub-column of  $R$ . We next prove a related claim concerning rows.

**Lemma 11** *For every sub-row  $L$  in  $R$ , such that  $L$  is assigned  $\overline{sum}(L)$  as a designated sum at some iteration of Procedure 2, we have that  $\widetilde{sum}(L) = \overline{sum}(L)$ .*

Observe that in particular we get that for every maximal sub-row  $L$  of  $R$ ,  $\widetilde{sum}(L) = \overline{sum}(L) = sum(L)$ .

**Proof:** Let  $\mathcal{L}$  the the set of sub-rows  $L$  of  $R$ , such that  $L$  is assigned  $\overline{sum}(L)$  as a designated sum at some iteration of Procedure 2. Observe that the set  $\mathcal{L}$  consists exactly of those rows that are maximal sub-rows for some  $R_p$ . We prove the claim by induction on the length of  $L \in \mathcal{L}$ . For the base of the induction, consider any sub-row  $L \in \mathcal{L}$  that is shortest among all sub-rows in  $\mathcal{L}$ . Since  $L$  is shortest, it must be completely filled in a single iteration as part of a block  $B$  (or otherwise there would be a shorter  $L' \subset L$  with a designated sum  $\overline{sum}(L')$ ). But by definition of the procedure, we get that  $\widetilde{sum}(L) = \overline{sum}(L)$  as required.

Assume that the claim holds for every  $L$  of length less than  $\ell$ , we prove it for  $L$  having length  $\ell$ . Consider the first iteration after which  $L$  became a maximal sub-row (and so received the designated sum  $\overline{sum}(L)$ ) in which part of  $L$  is filled. If all of  $L$  is filled, then the induction claim follows as in the base case. Otherwise, let  $x$  be the sum of elements that was filled in the part  $P \subset L$ . Let  $L'$  and  $L''$  be what remains of  $L$  to the left and right of  $P$  respectively. Then the procedure sets  $\overline{sum}(L') + \overline{sum}(L'') = \overline{sum}(L) - x$ . But  $L'$  and  $L''$  are strictly shorter than  $L$ , and therefore by the induction hypothesis  $\widetilde{sum}(L') = \overline{sum}(L')$  and  $\widetilde{sum}(L'') = \overline{sum}(L'')$ . Thus  $\widetilde{sum}(L) = \widetilde{sum}(L') + \widetilde{sum}(L'') + x = \overline{sum}(L') + \overline{sum}(L'') + x = \overline{sum}(L)$  as required. ■

**Definition 17 (Boundary)** We say that a point  $(i, j)$  is on the boundary of  $R$  if  $(i, j) \in R$ , but either  $(i + 1, j) \notin R$ , or  $(i, j + 1) \notin R$ , or  $(i + 1, j + 1) \notin R$ . We denote the set of boundary points by  $\mathcal{B}$ .

**Definition 18** For a point  $(i, j)$ ,  $1 \leq i, j \leq n$  let  $R^{\leq}(i, j)$  denote the subset of points  $(i', j') \in R$ ,  $i' \leq i, j' \leq j$ , and let  $sum^R(i, j) = \sum_{(i', j') \in R^{\leq}(i, j)} c_{i', j'}$  and  $\widetilde{sum}^R(i, j) = \sum_{(i', j') \in R^{\leq}(i, j)} \tilde{c}_{i', j'}$ .

Property 1 and therefore Theorem 3 will follow directly from the next two lemmas.

**Lemma 12** For every point  $(i, j) \in \mathcal{B}$ ,  $\widetilde{sum}^R(i, j) = sum^R(i, j)$ .

**Proof:** Consider any point  $(i, j) \in \mathcal{B}$  and let  $U = R^{\leq}(i, j)$ . Let  $\mathcal{C}(U) = \{B_1, \dots, B_q\}$  be the minimal set of blocks whose union contains  $U$ . For each  $B_h \in \mathcal{C}(U)$  we know that  $\widetilde{sum}(B_h) = sum(B_h)$ . In particular this is true for every  $B_h \subset U$ . Let  $\mathcal{C}_1(U) = \{B_h \in \mathcal{C}(U) : B_h \subset U\}$ . Hence we have that

$$\sum_{B_h \in \mathcal{C}_1(U)} \widetilde{sum}(B_h \cap U) = \sum_{B_h \in \mathcal{C}_1(U)} \widetilde{sum}(B_h) = \sum_{B_h \in \mathcal{C}_1(U)} sum(B_h). \quad (10)$$

If every  $B_h \in \mathcal{C}(U)$  is fully contained in  $U$  then  $\mathcal{C}_1(U) = \mathcal{C}(U)$  and we are done.

Otherwise, consider the remaining  $B_h$ 's in  $\mathcal{C}(U) \setminus \mathcal{C}_1(U)$  (i.e., blocks that are not fully contained in  $U$  but rather intersect it). Each of them either contains a column that is a sub-column of column  $j + 1$ , or a row that is a sub-row of row  $i + 1$ . Let the former subset be denoted  $\mathcal{C}_2(U)$  and the latter  $\mathcal{C}_3(U)$ . Thus  $\mathcal{C}_2(U)$  contains blocks that “intersect  $U$  from the right”, and  $\mathcal{C}_3(U)$  contain blocks that “intersect  $U$  from the top”. See for example Figure 6.

It is important to note that  $\mathcal{C}_2(U) \cap \mathcal{C}_3(U) = \emptyset$ : If there existed a block  $B_h \in \mathcal{C}_2(U) \cap \mathcal{C}_3(U)$ , it would necessarily contain both  $(i, j)$ , and the three neighboring points,  $(i + 1, j)$ ,  $(i, j + 1)$  and  $(i + 1, j + 1)$ . But this contradicts the fact that  $(i, j)$  is a boundary point.

For each  $B_h \in \mathcal{C}_2(U)$ ,  $B_h \cap U$  is a subset of maximal legal sub-columns with respect to  $R$  (since each  $B_h \in \mathcal{C}_2(U)$  cannot extend beyond row  $i$ ). Let us denote by  $\mathcal{K}_2(U)$  the set of all maximal legal




Figure 6: An illustration for the proof of Lemma 12. The solid line denotes the outline of  $U = R^{\leq}(i, j)$ , where point  $(i, j)$  is in the top-right corner. Blocks  $B_1$ – $B_6$  are fully contained in  $U$  and therefore belong to  $\mathcal{C}_1(U)$ . Blocks  $B_7$  and  $B_8$  belong to  $\mathcal{C}_2(U)$  and blocks  $B_9$ – $B_{11}$  belong to  $\mathcal{C}_3(U)$ . Block  $B_{10}$  is twice the size of  $B_9$  and  $B_{11}$  and so “extends out of the figure”.

sub-columns that belong to  $\bigcup_{B_h \in \mathcal{C}_2(U)} (B_h \cap U)$ . Since for every maximal legal sub-column  $K$ , we know that  $\widetilde{sum}(K) = sum(K)$ , we have that

$$\sum_{B_h \in \mathcal{C}_2(U)} \widetilde{sum}(B_h \cap U) = \sum_{K \in \mathcal{K}_2(U)} \widetilde{sum}(K) = \sum_{K \in \mathcal{K}_2(U)} sum(K). \quad (11)$$

Next consider the blocks  $B_h \in \mathcal{C}_3(U)$ . Let  $\mathcal{L}_3(U)$  be the set of sub-rows in  $U$  that are maximal sub-rows with respect to  $\bigcup_{B_h \in \mathcal{C}_3(U)} (B_h \cap U)$ . Thus,  $\bigcup_{B_h \in \mathcal{C}_3(U)} (B_h \cap U) = \bigcup_{L \in \mathcal{L}_3} L$ . We next observe that for every  $B_h \in \mathcal{C}_3(U)$ , all blocks that border  $B_h$  and belong either to  $\mathcal{C}_1(U)$  or to  $\mathcal{C}_2(U)$ , must be strictly smaller than  $B_h$ . This follows from the definition of legal sub-columns. Hence, the blocks in  $\mathcal{C}_1(U)$  and  $\mathcal{C}_2(U)$  are all removed before the blocks in  $\mathcal{C}_3(U)$ .

For each sub-row in  $\mathcal{L}_3(U)$  there exists the first iteration  $p$  in which it becomes a maximal sub-row with respect to  $R_p$  (following the removal of some block in  $\mathcal{C}_1(U) \cup \mathcal{C}_2(U)$  from  $R_{p-1}$ ). We partition the rows in  $\mathcal{L}_3(U)$  accordingly. That is, let  $\mathcal{L}_3^p(U)$  denote all sub-rows in  $\mathcal{L}_3(U)$ , that are maximal sub-rows with respect to  $R_p$  but were not maximal sub-rows with respect to  $R_{p-1}$ . Observe that in particular,  $\mathcal{L}_3^1(U)$  is the set of sub-rows in  $\mathcal{L}_3(U)$  that were already maximal sub-rows with respect to  $R$ . By this definition, necessarily the sub-rows in  $\mathcal{L}_3^p(U)$  constitute a sub-matrix of height  $s_{p-1}$ . By the second part of Lemma 10,  $\sum_{L \in \mathcal{L}_3^p(U)} \widetilde{sum}(L) = \sum_{L \in \mathcal{L}_3^p(U)} sum(L)$ , and by applying Lemma 11 we get that  $\sum_{L \in \mathcal{L}_3(U)} \widetilde{sum}(L) = \sum_{L \in \mathcal{L}_3(U)} sum(L)$ . Therefore,

$$\sum_{B_h \in \mathcal{C}_3(U)} \widetilde{sum}(B_h \cap U) = \sum_{L \in \mathcal{L}_3(U)} \widetilde{sum}(L) = \sum_{L \in \mathcal{L}_3(U)} sum(L). \quad (12)$$

By combining Equations (10)–(12) we get

$$\begin{aligned}
\widetilde{sum}(U) &= \sum_{B_h \in \mathcal{C}(U)} \widetilde{sum}(B_h \cap U) \\
&= \sum_{q=1}^3 \sum_{B_h \in \mathcal{C}_q(U)} \widetilde{sum}(B_h \cap U) \\
&= \sum_{B_h \in \mathcal{C}_1(U)} sum(B_h) + \sum_{K \in \mathcal{K}_2(U)} sum(K) + \sum_{L \in \mathcal{L}_3(U)} sum(L) \\
&= sum(U)
\end{aligned}$$

■




Figure 7: An illustration for the proof of Lemma 13.

**Lemma 13** *Let  $(i, j)$  be any point such that  $(i, j) \notin R$ . Then  $\widetilde{sum}^R(i, j) = sum^R(i, j)$ .*

**Proof:** Let  $(i', j') \in R$ ,  $i' < i$ ,  $j' \leq j$ , be the point for which  $j'$  is maximized, and if there are several such points, let it be the one amongst them for which  $i'$  is maximized. Thus,  $(i', j')$  is maximal in the sense that for every  $(i'', j'')$ ,  $i'' < i$ ,  $j'' \leq j$  such that  $(i'', j'') > (i', j')$  it holds that  $(i'', j'') \notin R$ . Furthermore, among all such maximal points it is the right-most one (i.e., it belongs to the column with the highest index). By definition,  $(i', j')$  belongs to  $\mathcal{B}$ , since  $(i' + 1, j')$  necessarily does not belong to  $R$ . Let  $\mathcal{L}(i, i', j)$  be the subset of all maximal sub-rows of  $R$  that belong to rows  $i' + 1, \dots, i$ , and end by column  $j$ . Then  $\widetilde{sum}^R(i, j) = \widetilde{sum}^R(i', j') + \sum_{L \in \mathcal{L}(i, i', j)} \widetilde{sum}(L)$ . By applying Lemma 12 and Lemma 11, we get that  $\widetilde{sum}^R(i, j) = sum^R(i, j)$ . ■

## 5.5 Distribution Matrices

As noted in the introduction, a sub-family of inverse Monge matrices that is of particular interest is the class of *distribution matrices*. A matrix  $V = \{v_{i,j}\}$  is said to be a distribution matrix, if



there exists a non-negative *density matrix*  $D = \{d_{i,j}\}$ , such that every entry  $v_{i,j}$  in  $V$  is of the form  $v_{i,j} = \sum_{k < i} \sum_{\ell < j} d_{k,\ell}$ . In particular, if  $V$  is a distribution matrix then the corresponding density matrix  $D$  is simply the matrix  $C'_V$  (as defined in Section 3). Hence, in order to test that  $V$  is a distribution matrix, we simply run our algorithm for inverse Monge matrix on  $C'_V$  instead of  $C_V$ .

## Acknowledgments

We would like to thank Noam Nisan for suggesting to examine combinatorial auctions in the context of property testing.

## References

- [BKR96] R.E. Burkard, B. Klinz, and R. Rudolf. Perspectives of monge properties in optimization. *Discrete Applied Mathematics and Combinatorial Operations Research and Computer Science*, 70:95–161, 1996.
- [BRW99] T. Batu, R. Rubinfeld, and P. White. Fast approximate pcps for multidimensional bin-packing problems. In *Proceedings of RANDOM*, pages 245–256, 1999.
- [DGL<sup>+</sup>99] Y. Dodis, O. Goldreich, E. Lehman, S. Raskhodnikova, D. Ron, and A. Samorodnitsky. Improved testing algorithms for monotonicity. In *Proceedings of RANDOM*, pages 97–108, 1999.
- [dVV00] S. de Vries and R. Vohra. Combinatorial auctions: a survey. available from: <http://www.kellogg.nwu.edu/faculty/vohra/htm/res.htm>, 2000.
- [EKK<sup>+</sup>00] F. Ergun, S. Kannan, S. R. Kumar, R. Rubinfeld, and M. Viswanathan. Spot-checkers. *JCSS*, 60(3):717–751, 2000.
- [Fis01] E. Fischer. The art of uninformed decisions: A primer to property testing. *Bulletin of the European Association for Theoretical Computer Science*, 75:97–126, 2001.
- [GGR98] O. Goldreich, S. Goldwasser, and D. Ron. Property testing and its connection to learning and approximation. *JACM*, 45(4):653–750, 1998.
- [GLS81] M. Grotschel, L. Lovasz, and A. Schrijver. The ellipsoid method and its consequences in combinatorial optimization. *Combinatorica*, 1:169–197, 1981.
- [Hof63] A.J. Hoffman. On simple linear programming problems. In *In Proceedings of Symposia in Pure Mathematics, Convexity*, volume 7, pages 317–327, 1963. American Mathematical Society.
- [IFF00] S. Iwata, L. Fleischer, and S. Fujishige. A combinatorial strongly polynomial algorithm for minimizing submodular functions. In *Proceedings of the Thirty-Fourth Annual ACM Symposium on the Theory of Computing*, pages 96–107, 2000. To appear in JACM.
- [LLN01] B. Lehmann, D. Lehmann, and N. Nisan. Combinatorial auctions with decreasing marginal utilities. In *ACM Conference on Electronic Commerce*, 2001.
- [Lov83] L. Lovász. Submodular functions and convexity. *Mathematical Programming: The State of the Art*, pages 235–257, 1983.

- [Ron01] D. Ron. Property testing. In *Handbook on Randomization, Volume II*, pages 597–649, 2001.
- [RS96] R. Rubinfeld and M. Sudan. Robust characterization of polynomials with applications to program testing. *SIAM Journal on Computing*, 25(2):252–271, 1996.
- [Sch00] A. Schrijver. A combinatorial algorithm minimizing submodular functions in strongly polynomial time. *Journal of Combinatorial Theory B*, 80:346–355, 2000.